



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
 International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁵ : C12N 15/11, C07H 21/00 A61K 31/70</p>	<p>A1</p>	<p>(11) International Publication Number: WO 94/01550 (43) International Publication Date: 20 January 1994 (20.01.94)</p>
<p>(21) International Application Number: PCT/US93/06326 (22) International Filing Date: 2 July 1993 (02.07.93) (30) Priority data: 07/909,069 2 July 1992 (02.07.92) US (60) Parent Application or Grant (63) Related by Continuation US 07/909,069 (CIP) Filed on 2 July 1992 (02.07.92) (71) Applicant (for all designated States except US): HYBRID- ON, INC. [US/US]; One Innovation Drive, Massachu- setts Biotechnology Research Park, Worcester, MA 01605 (US).</p>		<p>(72) Inventors; and (75) Inventors/Applicants (for US only) : AGRAWAL, Sudhir [IN/US]; 46G Shrewsbury Green Drive, Shrewsbury, MA 01545 (US). TANG, Jin-Yan [CN/US]; 16 Wells Street, #2L, Worcester, MA 06104 (US). (74) Agent: McDONNELL, John, J.; Allegretti & Witcoff, Ltd., Ten South Wacker Drive, Chicago, IL 60606 (US). (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI pa- tent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i></p>
<p>(54) Title: SELF-STABILIZED OLIGONUCLEOTIDES AS THERAPEUTIC AGENTS</p>		
<p>(57) Abstract</p> <p>The invention provides improved antisense oligonucleotides that are resistant to nucleolytic degradation. Such oligonucleotides are called self-stabilized oligonucleotides and comprise two regions: a target hybridizing region having a nucleotide sequence complementary to a nucleic acid sequence that is from a virus, a pathogenic organism, or a cellular gene; and a self-complementary region having an oligonucleotide sequence complementary to a nucleic acid sequence that is within the self-stabilized oligonucleotide.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic			SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TC	Togo
CZ	Czech Republic	MC	Monaco	UA	Ukraine
DE	Germany	MG	Madagascar	US	United States of America
DK	Denmark	ML	Mali	UZ	Uzbekistan
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

SELF-STABILIZED OLIGONUCLEOTIDES AS THERAPEUTIC AGENTS.

BACKGROUND OF THE INVENTION1. Field of the Invention

5 The invention relates to novel therapeutic agents used in the antisense oligonucleotide therapeutic approach. More particularly, the invention relates to improved antisense oligonucleotides that have increased resistance to nucleases.

10 2. Summary of the Related Art

 The antisense oligonucleotide based therapeutic principle provides an attractive strategy for rationally designing antiviral drugs and chemotherapeutic agents against other pathogens, and against disease conditions resulting from disorders of gene expression. The therapeutic principle relies upon specific binding between a target nucleic acid sequence and a complementary oligonucleotide. Several publications have demonstrated the efficacy of complimentary oligonucleotides in inhibiting gene expression by such specific binding.

 Zamecnik and Stephenson, Proc. Natl. Acad. Sci. USA 75: 285-288 (1978) discloses specific inhibition of Rous Sarcoma Virus replication in infected chicken fibroblasts by a 13-mer synthetic oligodeoxynucleotide that is complementary to part of the viral genome.

 Zamecnik et al., Proc. Natl. Acad. Sci. USA 83: 4143-4146 (1986) discloses inhibition of replication and expression of human immunodeficiency virus (HIV-1, then called HTLV-III) in cultured cells by synthetic oligonucleotide phosphodiester complementary to viral RNA.

-2-

More recently, it has been reported that oligonucleotides having greater resistance to nucleolytic degradation than oligonucleotide phosphodiester are more effective as antisense oligonucleotides. Agrawal, Tibtech 10: 152-158 (1992) has extensively reviewed the use of modified oligonucleotides as antiviral agents.

Sarin et al., Proc. Natl. Acad. Sci. USA 85: 7448-7451 (1988) teaches that oligodeoxynucleoside methylphosphonates are more active as inhibitors of HIV-1 than conventional oligodeoxynucleotides.

Agrawal et al., Proc. Natl. Acad. Sci. USA 85: 7079-7083 (1988) teaches that oligonucleotide phosphorothioates and various oligonucleotide phosphoramidates are more effective at inhibiting HIV-1 than conventional oligodeoxynucleotides.

Agrawal et al., Proc. Natl. Acad. Sci. USA 86: 7790-7794 (1989) discloses the advantage of oligonucleotide phosphorothioates in inhibiting HIV-1 in early and chronically infected cells.

An additional characteristic that renders oligonucleotides more effective as antisense agents is the ability to activate RNase H. Thus, oligonucleotide phosphorothioates, which are both resistant to nucleolytic degradation and activators of RNase H, are effective as inhibitors of HIV-1 and several other viruses.

Gao et al. Antimicrob. Agents and Chem. 34: 808 (1990) discloses inhibition of HSV by oligonucleotide phosphorothioates.

Storey et al., Nucleic Acids Res. 19: 4109 (1991) discloses inhibition of HPV by oligonucleotide phosphorothioates.

Leiter et al., Proc. Natl. Acad. Sci. USA 87: 3430 (1990) discloses inhibition of influenza virus by oligonucleotide phosphorothioates.

-3-

Unfortunately, oligonucleotide phosphorothioates increase resistance to nucleolytic degradation but do not provide complete resistance in vivo.

5 Agrawal et al., Proc. Natl. Acad. Sci. USA 88: 7595-7599 (1991) teaches that oligonucleotide phosphorothioates are extensively degraded from the 3' end in mice.

10 In addition, oligonucleotide phosphorothioates form less stable duplexes between the oligonucleotide and target than oligodeoxynucleotides phosphodiester. To overcome these deficiencies, oligonucleotides having cap structures at the 3' terminus have been developed. Agrawal and Goodchild, Tetrahedron Lett. 28: 3539-3542 (1987) discloses the use of oligodeoxynucleoside 15 methylphosphonates as 5' and 3' capping agents. Shaw et al., Nucleic Acids Res. 19: 747-750 (1991) discloses oligodeoxynucleotide phosphodiester having blocking structures at the 3' end.

20 Temsamani et al., in Antisense Strategies, Annals of New York Academy of Sciences (in press) (1992) discloses 3' capped oligonucleotide phosphorothioates.

25 Even these nuclease resistant 3' capped oligonucleotides can become degraded eventually as the 3' capped end of these oligonucleotides is slowly digested by a combination of endonuclease and exonuclease activities.

30 There is, therefore, a need for oligonucleotides that form stable duplexes, resist nucleolytic degradation and activate RNase H, without the disadvantages of oligonucleotides that are known in the art. Ideally, such oligonucleotides should resist even the combined effect of endonucleases and exonucleases, should stably pair with target sequences at physiological temperatures,

-4-

should activate RNase H and should produce only nucleosides as degradation products.

5 Oligonucleotides having self-complementary structures that can result in hairpin formation are known in the art.

Germann et al., *Biochemistry* 24: 5698-5702 (1985) discloses a partially self-complementary 24-mer oligonucleotide, d(GC)₅T₄(CG)₅, that undergoes a β -DNA to Z-DNA transition.

10 Hilbers et al., *Biochimie* 67: 685-695 (1985) discusses the dynamics of hairpin formation in a partially self-complementary oligonucleotide, dATCCTAT_nTAGGAT.

15 Neither of these physical studies related to either oligonucleotide stability or to therapeutic use of oligonucleotides.

20 Thus, the prior art is devoid of any teaching or suggestion about using self-complementary oligonucleotides in the antisense oligonucleotide therapeutic approach, nor does it discuss the use of hairpin formation as a means of rendering an oligonucleotide resistant to nucleolytic degradation.

-5-

BRIEF SUMMARY OF THE INVENTION

The invention relates to novel therapeutic agents used in the antisense oligonucleotide therapeutic approach. The invention provides improved antisense oligonucleotides that are resistant to nucleolytic degradation. Oligonucleotides according to the invention resist nucleolytic degradation, including the combined effect of endonucleases and exonucleases. Oligonucleotides according to the invention form stable hybrids with target sequences under physiological conditions, activate RNase H and produce only nucleosides as degradation products.

The advantages of oligonucleotides according to the invention, known as self-stabilized oligonucleotides, arise from the presence of two structural features: a target hybridizing region and a self-complementary region. The target hybridizing region comprises an oligonucleotide sequence that is complementary to a nucleic acid sequence that is from a plant or animal virus, a pathogenic organism, or a cellular gene or gene transcript, the abnormal gene expression or product of which results in a disease state. The self-complementary region comprises an oligonucleotide sequence that is complementary to a nucleic acid sequence within the oligonucleotide. Thus, at least when the oligonucleotide is not hybridized to a target nucleic acid sequence, the oligonucleotide forms a totally or partially double-stranded structure that is resistant to nucleolytic degradation. Since the inherent structure of these molecules confers resistance to nucleases, it is not necessary to use modified internucleotide linkages to confer such resistance, although of course, modified linkages may be used. Thus, the use of oligonucleotide phosphodiester or oligonucleotide phosphorothioates, both of which are degraded in vivo, is made feasible by oligonucleotides according to the invention. This

-6-

results in oligonucleotides that activate RNase H, an important feature for the antisense therapeutic compound. Also, the use of oligonucleotide phosphodiester provides more stable hybridization between therapeutic oligonucleotides and target sequences. Finally, degradation of such oligonucleotides results only in nucleotide breakdown products, thus minimizing potential toxicity. These advantages result in a superior therapeutic oligonucleotide.

10 The invention further provides self-stabilized ribozymes, since the self-complementary motif of the invention can be conveniently used with ribonucleotides. Such ribozymes according to the invention have generally typical ribozyme structure, except that they have a self-complementary region at or near the 5' or 3' end. This region confers nuclease resistance upon the ribozymes, making them more stable than ribozymes that are known in the art.

-7-

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates a self-stabilized oligonucleotide according to the invention in hairpin and hybridized configurations.

5 Figure 2 illustrates a self-stabilized oligonucleotide according to the invention in hammer-like configuration.

10 Figure 3 shows results of duplex stability studies for hybridization between oligonucleotides or self-stabilized oligonucleotides and complementary target oligonucleotides.

Figure 4 shows results of 3'-exonuclease treatment of oligonucleotides.

15 Figure 5 shows the structure of self-stabilized oligonucleotides used in Examples 1-4.

Figure 6 shows a mechanism of therapeutic action of self-stabilized oligonucleotides.

20 Figure 7 shows a self-stabilized ribozyme according to the invention. This example of a self-stabilized ribozyme according to the invention is complementary to the HIV gag region and results in the cleavage of a HIV gag mRNA.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The invention relates to novel therapeutic agents that are useful in the treatment of virus infections, infections by pathogenic organisms, and diseases arising
5 from abnormal gene expression or gene products.

In a first aspect, the invention provides therapeutic self-stabilized oligonucleotides that are more resistant to nucleolytic degradation than oligonucleotides that are known in the art. For purposes
10 of the invention, the term oligonucleotide includes polymers of ribonucleotides, deoxyribonucleotides, or both, with ribonucleotide and/or deoxyribonucleotide monomers being connected together via 5' to 3' linkages which may include any of the linkages that are known in
15 the antisense oligonucleotide art. In addition, the term oligonucleotides includes such molecules having modified nucleic acid bases and/or sugars, as well as such molecules having added substituents, such as diamines, cholesteryl or other lipophilic groups. Certain
20 preferred combinations of monomers and inter-monomer linkages are discussed in greater detail below.

Oligonucleotides according to the invention are generally characterized by having two regions: a target hybridizing region and a self-complementary region. A
25 first embodiment of a self-stabilized oligonucleotide according to the invention is shown in Figure 1. In this embodiment, the target hybridizing region is shown as connected rectangular squares, and the self-complementary region is shown as connected circles. The complementary
30 nucleic acid sequence in a target RNA molecule is represented by connected diamonds. Hydrogen bonding between nucleotides is indicated by dots. The oligonucleotide is stabilized, i.e., rendered resistant to nucleolytic degradation from the 5' or 3' end by base-
35 pairing between the target hybridizing region and the

-9-

self-complementary regions and/or by base-pairing between complementary sequences within the self-complementary region. When the oligonucleotide encounters a nucleic acid molecule having a complementary nucleic acid sequence, base-pairing between the target hybridizing region and the self-complementary region of the oligonucleotide is disrupted and replaced by base-pairing between the target hybridizing region of the oligonucleotide and the complementary nucleic acid sequence of the target nucleic acid molecule. This disruption and replacement of base-pairing takes place because the intermolecular base-paired structure formed by the hybrid between the target nucleic acid sequence and the target hybridizing region is more thermodynamically stable than the intra-molecular base-paired structure formed by the self-complementary oligonucleotide. This phenomenon is illustrated in Figure 3 and discussed in greater detail in Example 4.

A second embodiment of an oligonucleotide according to the invention operates in a similar way as the first embodiment, but forms a different structure upon self-complementary base-pairing. This alternative embodiment forms a hammer-like structure as shown in Figure 2. In this embodiment, the self-complementary region contains oligonucleotide sequences that can base pair with other oligonucleotide sequences within the self-complementary region. The self complementary region may also contain oligonucleotide sequences that are complementary to the target hybridizing region.

The target hybridizing region of an oligonucleotide according to the invention has an oligonucleotide sequence that is complementary to a nucleic acid sequence that is from a virus, a pathogenic organism, or a cellular gene or gene transcript, the abnormal gene expression or product of which results in a disease state. Preferably the target hybridizing region is from

-10-

about 8 to about 50 nucleotides in length. For purposes of the invention, the term "oligonucleotide sequence that is complementary to a nucleic acid sequence" is intended to mean an oligonucleotide sequence (2 to about 50 nucleotides) that hybridizes to the nucleic acid sequence under physiological conditions, e.g., by Watson-Crick base pairing (interaction between oligonucleotide and single-stranded nucleic acid) or by Hoogsteen base pairing. (interaction between oligonucleotide and double-stranded nucleic acid) or by any other means. Such hybridization under physiological conditions is measured as a practical matter by observing interference with the function of the nucleic acid sequence.

The nucleic acid sequence to which the target hybridizing region of an oligonucleotide according to the invention is complementary will vary, depending upon the disease condition to be treated. In many cases the nucleic acid sequence will be a virus nucleic acid sequence. The use of antisense oligonucleotides to inhibit various viruses is well known, and has recently been reviewed in Agrawal, Tibtech 10:152-158 (1992). Viral nucleic acid sequences that are complementary to effective antisense oligonucleotides have been described for many viruses, including human immunodeficiency virus type 1 (U.S. Patent No. 4,806,463, the teachings of which are herein incorporated by reference), Herpes simplex virus (U.S. Patent No. 4,689,320, the teachings of which are hereby incorporated by reference), Influenza virus (U.S. Patent No. 5,XXX,XXX; Ser. No. 07/516,275, allowed June 30, 1992; the teachings of which are hereby incorporated by reference), and Human papilloma virus (Storey et al., Nucleic Acids Res. 19:4109-4114 (1991)). Sequences complementary to any of these nucleic acid sequences can be used for the target hybridizing region of oligonucleotides according to the invention, as can be oligonucleotide sequences

-11-

complementary to nucleic acid sequences from any other virus. Additional viruses that have known nucleic acid sequences against which antisense oligonucleotides can be prepared include Foot and Mouth Disease Virus (See 5 Robertson et al., J. Virology 54: 651 (1985); Harris et al., J. Virology 36: 659 (1980)), Yellow Fever Virus (See Rice et al., Science 229: 726 (1985)), Varicella-Zoster Virus (See Davison and Scott, J. Gen. Virology 67: 2279 (1986), and Cucumber Mosaic Virus (See Richards et al., 10 Virology 89: 395 (1978)).

Alternatively, the target hybridizing region of oligonucleotides according to the invention can have an oligonucleotide sequence complementary to a nucleic acid sequence of a pathogenic organism. The nucleic acid 15 sequences of many pathogenic organisms have been described, including the malaria organism, Plasmodium falciparum, and many pathogenic bacteria. Oligonucleotide sequences complementary to nucleic acid sequences from any such pathogenic organism can form the 20 target hybridizing region of oligonucleotides according to the invention. Examples of pathogenic eukaryotes having known nucleic acid sequences against which antisense oligonucleotides can be prepared include Trypanosoma brucei gambiense and Leishmania (See Campbell 25 et al., Nature 311: 350 (1984)), Fasciola hepatica (See Zurita et al., Proc. Natl. Acad. Sci. USA 84: 2340 (1987)). Antifungal oligonucleotides can be prepared using a target hybridizing region having an oligonucleotide sequence that is complementary to a 30 nucleic acid sequence from, e.g., the chitin synthetase gene, and antibacterial oligonucleotides can be prepared using, e.g., the alanine racemase gene.

In yet another embodiment, the target hybridizing region of oligonucleotides according to the invention can 35 have an oligonucleotide sequence complementary to a

-12-

cellular gene or gene transcript, the abnormal expression or product of which results in a disease state. The nucleic acid sequences of several such cellular genes have been described, including prion protein (Stahl and Prusiner, FASEB J. 5: 2799-2807 (1991)), the amyloid-like protein associated with Alzheimer's disease (U.S. Patent No. 5,015,570, the teachings of which are hereby incorporated by reference), and various well-known oncogenes and proto-oncogenes, such as c-myb, c-myc, c-abl, and n-ras. In addition, oligonucleotides that inhibit the synthesis of structural proteins or enzymes involved largely or exclusively in spermatogenesis, sperm motility, the binding of the sperm to the egg or any other step affecting sperm viability may be used as contraceptives for men. Similarly, contraceptives for women may be oligonucleotides that inhibit proteins or enzymes involved in ovulation, fertilization, implantation or in the biosynthesis of hormones involved in those processes.

Hypertension can be controlled by oligodeoxynucleotides that suppress the synthesis of angiotensin converting enzyme or related enzymes in the renin/angiotensin system; platelet aggregation can be controlled by suppression of the synthesis of enzymes necessary for the synthesis of thromboxane A2 for use in myocardial and cerebral circulatory disorders, infarcts, arteriosclerosis, embolism and thrombosis; deposition of cholesterol in arterial wall can be inhibited by suppression of the synthesis of fattyacyl co-enzyme A: cholesterol acyl transferase in arteriosclerosis; inhibition of the synthesis of cholinephosphotransferase may be useful in hypolipidemia.

There are numerous neural disorders in which hybridization arrest can be used to reduce or eliminate adverse effects of the disorder. For example, suppression of the synthesis of monoamine oxidase can be

-13-

used in Parkinson's disease; suppression of catechol o-methyl transferase can be used to treat depression; and suppression of indole N-methyl transferase can be used in treating schizophrenia.

5 Suppression of selected enzymes in the arachidonic acid cascade which leads to prostaglandins and leukotrienes may be useful in the control of platelet aggregation, allergy, inflammation, pain and asthma.

10 Suppression of the protein expressed by the multidrug resistance (mdr) gene, which is responsible for development of resistance to a variety of anti-cancer drugs and is a major impediment in chemotherapy may prove to be beneficial in the treatment of cancer.

15 Oligonucleotide sequences complementary to nucleic acid sequences from any of these genes can be used for the target hybridizing region of oligonucleotides according to the invention, as can be oligonucleotide sequences complementary to any other cellular gene or gene transcript, the abnormal expression or product of which results in a disease state.

20 Antisense regulation of gene expression in plant cells has been described in U.S. Patent No. 5,107,065, the teachings of which are hereby incorporated by reference.

25 In a second aspect, the invention provides nuclease resistant oligonucleotides that activate RNase H. The target hybridizing region of oligonucleotides according to the invention may contain ribonucleotides, deoxyribonucleotides or any analogs of ribonucleotides or deoxyribonucleotides. In one preferred embodiment, this region is composed of ribonucleotides. In another preferred embodiment, this region is composed of deoxyribonucleotides. In yet another preferred embodiment, this region comprises a mixture of ribonucleotides and deoxyribonucleotides. An additional

30

35

-14-

preferred embodiment has a target hybridizing region comprising oligonucleotide phosphodiester, phosphorothioate, or phosphorodithioate, or mixtures or these with ribonucleotides or deoxyribonucleotides. 5 These preferred embodiments all provide for the activation of RNase H, as long as four or more contiguous deoxyribonucleotide phosphodiester, phosphorothioate, or phosphorodithioate are present. Of course, other 10 embodiments employing target hybridizing regions that do not activate RNase H can also be made.

Synthesis procedures for each of these embodiments are well known in the art. Both oligodeoxyribonucleotide phosphodiester and oligodeoxyribonucleotide phosphorothioate and their analogs can be synthesized by 15 the H-phosphonate approach described in U.S. Patent No. 5, , , (Ser. No. 07/334,679; allowed on March 19, 1992), the teachings of which are hereby incorporated by reference. The H-phosphonate approach can also be used to synthesize oligoribonucleotides and 20 oligoribonucleotide analogs, as described in Agrawal and Tang, Tetrahedron Lett. 31: 7541-7544 (1990). Synthesis of oligonucleotide phosphorodithioate is also known in the art.

Of course, many other embodiments are possible, and 25 those skilled in the art will recognize that other analogs or combinations of analogs can be used in the target hybridizing region of oligonucleotides according to the invention. Such analogs are characterized by having internucleotide linkages other than the natural 30 phosphodiester linkage. The synthesis of many such analogs is well known in the art, including analogs having alkylphosphonate, (Agrawal and Goodchild, Tetrahedron Lett. 28: 3539-3542 (1987)) or phosphoramidate (Agrawal et al., Proc. Natl. Acad. Sci. 35 USA 85: 7079-7083 (1988)) linkages.

-15-

The second significant region of self-stabilized oligonucleotides according to the invention is the self-complementary region. The self-complementary region contains oligonucleotide sequences that are complementary to other oligonucleotide sequences within the oligonucleotide. These other oligonucleotide sequences may be within the target hybridizing region or within the self-complementary region, or they may span both regions. The complementary sequences form base pairs, resulting in the formation of a hairpin structure, as shown in Figure 1, or a hammer-like structure, as shown in Figure 2. Either the hairpin structure or the hammer-like structure can have loops resulting from non-base-paired nucleotides, as shown in Figure 1 for the hairpin structure, or can be devoid of such loops, as shown in Figure 2 for the hammer-like structure. The number of base-pairs to be formed by intra-molecular hybridization involving the self-complementary region may vary, but should be adequate to maintain a double-stranded structure so that the 3' end is not accessible to endonucleases. Generally, about 4 or more base-pairs will be necessary to maintain such a double-stranded structure. In a preferred embodiment, there are about 10 intramolecular base-pairs formed in the self-stabilized oligonucleotide, with the 10 base pairs being consecutive and involving the 3'-most nucleotides. Of course, the intra-molecular base-pairing can be so extensive as to involve every nucleotide of the oligonucleotide. Preferably, this will involve a self-complementary region of about 50 nucleotides or less.

In one preferred embodiment the self-complementary region may be connected to the target hybridizing region by a suitable non-nucleic acid linker. Examples of such linkers include substituted or unsubstituted alkyl groups. In one most preferred embodiment the linker is a (ethylene glycol)₁₋₆ linker. At the larger size for

-16-

this linker, the synthesis may be conveniently carried out by using commercially available triethylene glycol that has a dimethyltrityl protective group at one end and a cyanoethylphosphoramidite group at the other end.

5 The self-complementary region may contain
ribonucleotides, deoxyribonucleotides, analogs of
ribonucleotides or deoxyribonucleotides having artificial
linkages, or combinations of any of the above. The
10 ability to activate RNase H is not important for the
self-complementary region, so nucleotides having
artificial linkages that do not activate RNase H can be
used in this region without diminishing the effectiveness
of the oligonucleotide. Thus, in addition to
15 phosphodiester and phosphorothioate or phosphorodithioate
linkages, this region may also or alternatively contain
phosphoramidate (including N-substituted
phosphoramidates), alkylphosphonate,
alkylphosphonothioate linkages as well as non-phosphate
20 containing linkages, such as sulfone, sulfate, and keto
linkages. Of course, in non-RNase H activating
embodiments of self-stabilized oligonucleotides according
to the invention, any of these linkages can be used in
the target hybridizing region as well.

 In one preferred embodiment, the self-stabilized
25 oligonucleotide is rendered hyperstabilized. This may be
accomplished by incorporating into the self-complementary
region one or more ribonucleotides or 2'-O-Me-
ribonucleotides, wherein the complementary portion of the
target hybridizing region is DNA. Alternatively, the
30 complementary region of the target hybridizing region may
contain ribonucleotides or 2'-O-Me-ribonucleotides, and
the self-complementary region may contain DNA. These
oligonucleotides will be hyperstabilized because the
interaction between DNA and RNA is more stable than the
35 interaction between DNA and DNA. Yet another way in
which the self-complementary region (and/or the linker

-17-

region) may be modified to yield a hyperstabilized self-stabilized oligonucleotide is to incorporate one or more intercalating agent molecule. These oligonucleotides are hyperstabilized because the intercalating agent stabilizes the hybrid formed between the self-complementary region and the target hybridizing region. Any intercalating agent is acceptable for this purpose. Preferred intercalating agents include acridine and ethidium. Oligonucleotides containing acridine are readily prepared by using the commercially available acridine-ON phosphoramidite, or 3'-acridine-ON CPG (Clontech Laboratories, Inc.).

In a third aspect, the invention provides ribozymes that are more stable than ribozymes that are known in the art. Ribozymes are catalytic RNA molecules that cleave internucleotide bonds. The stability of such ribozymes according to the invention is provided by the incorporation of a self-complementary region at or near the 5' or 3' end of the ribozyme molecule. This self-complementary region results in the formation of a hairpin or hammer-like structure, thus rendering the 5' or 3' end of the molecule double-stranded, which causes the ribozyme molecule to resist nucleolytic degradation. The structure and function of ribozymes is generally taught in U.S. Patent No. 4,987,071, the teachings of which are hereby incorporated by reference.

In a fourth aspect, the invention provides a method for inhibiting the gene expression of a virus, a pathogenic organism or a cellular gene, the method comprising the step of providing self-stabilized oligonucleotides or ribozymes according to the invention to cells infected with the virus or pathogenic organism in the former two cases or to cells generally in the latter case.

-18-

5 In a fifth aspect, the invention provides a method
of treating a diseased human or animal in which the
disease results from infection with a virus or pathogenic
organism, or from the abnormal expression or produce of
a cellular gene. The method comprises administering
self-stabilized oligonucleotides according to the
invention in a pharmaceutically acceptable carrier to the
diseased human or animal. Preferably, the routes of such
administration will include oral, intranasal, rectal and
10 topical administration. In such methods of treatment
according to the invention the self-stabilized
oligonucleotides may be administered in conjunction with
other therapeutic agents, e.g., AZT in the case of AIDS.

15 A variety of viral diseases may be treated by the
method of treatment according to the invention, including
AIDS, ARC, oral or genital herpes, papilloma warts, flu,
foot and mouth disease, yellow fever, chicken pox,
shingles, HTLV-leukemia, and hepatitis. Among fungal
diseases treatable by the method of treatment according
20 to the invention are candidiasis, histoplasmosis,
cryptococcosis, blastomycosis, aspergillosis,
sporotrichosis, chromomycosis, dematophytosis and
coccidioidomycosis. The method can also be used to treat
rickettsial diseases (e.g., typhus, Rocky Mountain
25 spotted fever), as well as sexually transmitted diseases
caused by Chlamydia trachomatis or Lymphogranuloma
venereum. A variety of parasitic diseases can be treated
by the method according to the invention, including
amebiasis, Chagas' disease, toxoplasmosis,
30 pneumocystosis, giardiasis, cryptosporidiosis,
trichomoniasis, and Pneumocystis carini pneumonia; also
worm (helminthic diseases) such as ascariasis,
filariasis, trichinosis, schistosomiasis and nematode or
cestode infections. Malaria can be treated by the method
35 of treatment of the invention regardless of whether it is

-19-

caused by P. falciparum, P. vivax, P. orale, or P. malariae.

5 The infectious diseases identified above can all be treated by the method of treatment according to the invention because the infectious agents for these diseases are known and thus self-stabilized oligonucleotides according to the invention can be prepared, having a target hybridizing region that has an oligonucleotide sequence that is complementary to a
10 nucleic acid sequence that is an essential nucleic acid sequence for the propagation of the infectious agent, such as an essential gene.

15 Other disease states or conditions that are treatable by the method according to the invention result from an abnormal expression or product of a cellular gene. These conditions can be treated by administration of self-stabilized oligonucleotides according to the invention, and have been discussed earlier in this disclosure.

20 The invention provides numerous advantages over oligonucleotides that are known in the art. First, the self-stabilized oligonucleotides according to the invention have a longer half-life than most known oligonucleotides, thereby lowering the dosage that will
25 be required for therapeutic efficacy. Even greater resistance to nuclease degradation can be provided by using nuclease resistant internucleotide linkages near or cap structures at one or both ends of the oligonucleotide. Second, the enzymatic stability
30 afforded by the base-paired structures involving the self-complementary sequences allows the use of oligonucleotide phosphodiester, which otherwise are rapidly degraded. This provides the advantages of increased duplex stability and RNase H activation, which
35 are not both provided by any nuclease resistant

-20-

oligonucleotide known in the art. The advantage of RNase H activation is retained when oligonucleotide phosphorothioates or phosphorodithioates are used. A third advantage is that the only degradation product of several embodiments of oligonucleotides according to the invention is nucleotides, e.g., nucleoside monophosphates and/or nucleoside monothiophosphates. Finally, the invention allows the use of either deoxyribonucleosides or ribonucleosides. The ability to use the latter makes the invention readily adaptable for use with ribozymes, for which enzymatic stability is critical.

The following examples are provided to further illustrate certain aspects of preferred embodiments of the invention, and are not intended to be limiting in nature.

EXAMPLE 1

Nuclease Resistance of Oligonucleotide Phosphodiester

The oligonucleotides used in the study are shown in Figure 5. Oligonucleotide CMPD A is complementary to a portion of the gag region of HIV-1. Oligonucleotide CMPD B uses this same region as a target hybridizing region, but adds a 3' self-complementary region of 10 nucleotides. Oligonucleotides CMPD E and CMPD F are identical to CMPD B, except that the self-complementary regions of CMPD E and CMPD F are 6 and 4 nucleotides, respectively. Oligonucleotide CMPD G is identical to CMPD A, except that it has 10 mismatched nucleotides (T₁₀) added at its 3' end.

The oligonucleotides were tested for their relative resistance to 3' exonucleolytic degradation. For each oligonucleotide, 0.4 A₂₆₀ units of oligonucleotide was lyophilized, dissolved in 0.5ml buffer (10mM Tris, 10mM MgCl₂, pH 8.5) and mixed with 5μl (1.5 milliunits) of

-21-

snake venom phosphodiesterase (SVPD). The mixture was incubated at 37°C in a thermally regulated cell and A_{260} was plotted against time. Oligonucleotide degradation was measured as function of increase in hyperchromicity.

5 The results of these experiments are shown in Table I, below. These results demonstrate that self-stabilized oligonucleotide phosphodiesterases according to the invention are far more resistant to 3' exonucleolytic degradation than either oligonucleotide phosphodiesterases or oligonucleotide phosphodiesterases having a non-complementary tail.

10 In addition to the testing described above, the oligonucleotides were also subjected to DNA Polymerase I 3'-exonuclease digestion. As shown in Figure 4 the non-self-stabilized oligonucleotides, CMPDs A and G were digested to completion in 30 minutes, whereas self-stabilized CMPD B was only partly digested over 30 minutes.

20 TABLE I

HALF-LIFE OF OLIGONUCLEOTIDES

<u>Oligonucleotide</u>	<u>t_{1/2} for SVPD digestion</u>
CMPD A	75 seconds
CMPD G	75 seconds
CMPD B	950 seconds

25

EXAMPLE 2

Nuclease Resistance of Oligonucleotide Phosphorothioates

To test the relative nuclease resistance of self-stabilized and non-self-stabilized oligonucleotide phosphorothioates, a DNA Polymerase I 3'-exonuclease activity assay was used, because of the slow degradation of oligonucleotide phosphorothioates by SVPD.

30 All oligonucleotides were labelled at the 5'- end with gamma-³²P-ATP and kinase. To a solution of 40 pmole

-22-

5'-labelled oligonucleotide in 20 μ l buffer (40 mM Tris. HCl pH 8.0, 10 mM MgCl₂, 5 mM DTT, 50 mM KCl, 50 μ g/ml BSA), 5 units DNA polymerase I was added and incubated at 37°C. Aliquots of 4 μ l were taken at 0, 30, 60, 120
5 minutes and were mixed with 6 μ l stop solution (98% formamide, 10 mM EDTA, 0.1% xylene cyanol, 0.1% bromophenol blue). The samples were analyzed by 15% acrylamide gel (urea) and autoradiography.

The results are shown in Figure 4. Phosphorothioate analog of CMPD A was digested to almost 50% within 4
10 hours. The phosphorothioate analog of CMPD B, however, was undegraded after 4 hours. Phosphorothioate analogs of CMPD E and F, which have 6 and 4 base pairs of self-complementary sequence, respectively were also found to
15 be stable. Phosphorothioate analog of CMPD G, having extended structure, but no self-complementary region, was digested at same rate as CMPD A. These results demonstrate that self-stabilized oligonucleotide phosphorothioates are far more resistant to nucleolytic
20 degradation than are non-self-stabilized oligonucleotide phosphorothioates.

EXAMPLE 3

Anti-HIV Activity Of Oligonucleotides

Self-stabilized and non-self stabilized
25 oligonucleotide phosphodiester were tested for their ability to inhibit HIV-1 in tissue culture. The oligonucleotides used in this study are shown in Figure 5.

H9 lymphocytes were infected with HIV-1 virions
30 (\approx 0.01 - 0.1 TCID₅₀/cell) for one hour at 37°C. After one hour, unadsorbed virions were washed and the infected cells were divided among wells of 24 well plates. To the infected cells, an appropriate concentration (from stock solution) of oligonucleotide was added to obtain the
35 required concentration in 2 ml medium. In a positive

-23-

control experiment ddC or AZT was added. The cells were then cultured for three days. At the end of three days, supernatant from the infected culture was collected and measured for p24 expression by ELISA. The level of expression of p24 was compared between oligonucleotide treated and untreated (no drug) infected cells.

5

Cytotoxicity of oligonucleotides was studied by culturing the cells with increasing concentration of oligonucleotide and by the trypan blue dye exclusion method.

10

The results of two experiments are shown in Table III, below.

-24-

TABLE III

Anti-HIV Activity of OligonucleotidesExperiment 1

	Concentration (g/ml)	Inhibition of p24 (%)	% Cell Survival	IC ₅₀ (g/ml)
CMPD A	25	90	93	2
	5	89	103	
	1	15	94	
	0.2	26	97	
CMPD B	25	90	95	0.25
	5	85	92	
	1	84	94	
	0.2	46	103	
CMPD G	25	86	106	0.5
	5	86	105	
	1	81	106	
	0.2	0	109	
AZT	0.2	90	95	0.037 μM
	0.04	73	98	
	0.08	44	104	
	.0016	6	108	

5

-25-

Experiment 2

	Concentration (g/ml)	Inhibition of p24 (%)	% Cell Survival	IC ₅₀ (g/ml)
CMPD A	5	66	93	2.8
	1	20	101	
	0.2	21	107	
	0.04	0	102	
CMPD B	5	93	89	0.35
	1	81	99	
	0.2	33	103	
	0.04	0	104	
CMPD E	5	89	93	0.45
	1	41	100	
	0.2	19	99	
	0.04	0	102	
CMPD F	5	89	93	1.5
	1	41	100	
	0.2	19	99	
	0.04	0	102	
AZT	0.2	89	93	0.1 μ m
	0.04	65	98	
	0.008	5	101	
	0.0016	6	103	

5

10

All self-stabilized oligonucleotides exhibited greater anti-HIV activity than CMPD A, the non-self-stabilized oligonucleotide. Greatest activity was observed for the self-stabilized oligonucleotide having 10 self-complementary nucleotides, which exhibited nearly ten times the activity of the oligonucleotide phosphodiester. The oligonucleotide CMPD G, which has a poly T tail, also showed some increase in activity,

-26-

probably as a result of stabilization from hybridization with polyA from mRNA in the H9 cells.

5 The probable mechanism of action of the CMPD B oligonucleotide is shown in Figure 6. The oligonucleotide enters the cell in a partially double-stranded form as a result of intramolecular base-pairing involving the self-complementary region. As the oligonucleotide encounters on HIV RNA molecule having a nucleic acid sequence that is complementary to the
10 oligonucleotide sequence of the target hybridizing region of the oligonucleotide, hybridization occurs between the HIV RNA and the target hybridizing region. This hybridization disrupts the intramolecular hybridization involving the self-complementary region. RNase H
15 activity then cleaves the HIV RNA, allowing the oligonucleotide to once again self-stabilize by intramolecular base-pairing.

To test the relative anti-HIV activity of additional oligonucleotide structures, the above experiment was
20 repeated using additional oligonucleotides, as well as the oligonucleotides described in Experiments 1 & 2. The additional oligonucleotides are shown in Figure 5. These additional oligonucleotides were CMPD C, in which the self-complementary region is complementary to the
25 oligonucleotide through its 5' end; CMPD D, which has a 8 nucleotide self-complementary region; and CMPD H, a 35 mer oligonucleotide having perfect complementarity to the HIV gag RNA, but no self-complementary region. The results of this third experiment are shown in Table IV,
30 below.

These results demonstrate that fully self-complementary self-stabilized oligonucleotides are roughly equivalent in anti-HIV activity to partially self-complementary self-stabilized oligonucleotides. The
35 results also show that four self-complementary nucleotides are adequate to confer enhanced efficacy.

-27-

TABLE IV
Anti-HIV Activity of Oligonucleotides

Experiment 3

	Concentration (g/ml)	Inhibition of p24 (%)	% Cell Survival	IC ₅₀ (g/ml)
CMPD A	5.0	92	97	1.7
	1.0	36	103	
	0.2	23	102	
	0.04	0	109	
CMPD B	5.0	95 (97)*	98 (97)*	0.5 (0.2)*
	1.0	61 (74)*	101 (102)*	
	0.2	33 (49)*	104 (103)*	
	0.04	0 (19)*	11 (106)*	
CMPD G	5.0	94	97	0.6
	1.0	68	104	
	0.2	11	109	
	0.04	12	110	
CMPD E	5.0	92	98	0.8
	1.0	55	101	
	0.2	13	103	
	0.04	0	107	
CMPD F	5.0	95	99	0.25
	1.0	64	102	
	0.2	48	104	
	0.04	22	109	
CMPD C	5.0	94	96	0.3
	1.0	76	101	
	0.2	39	103	
	0.04	17	106	

*Results of second independent screening.

-28-

	Concentration ($\mu\text{g/ml}$)	Inhibition of p24 (%)	% Cell Survival	IC ₅₀ ($\mu\text{g/ml}$)
CMPD H	5	92	93	0.26
	1	88	101	
	.2	43	98	
	.04	0	102	
CMPD D	5	80	93	0.4
	1	76	100	
	.2	30	108	
	.04	3	109	

EXAMPLE 4

**Stability of Duplexes Between Self-Stabilized
Oligonucleotides and Complementary Oligos**

5

To test the stability of duplexes formed between self-stabilized oligonucleotides and complementary nucleic acid sequences hybridization studies were carried out. In a first study oligonucleotide CMPD A, which lacks self-complementary sequences, was mixed at room temperature with a complementary 25-mer oligonucleotide. The mixture was then gradually heated and increase in hyperchromicity was plotted against increase in temperature. In this study, the results of which are shown as a dotted line in Figure 3, the melting temperature of the duplex was found to be about 65°C.

10

15

In a second study CMPD B, having the same target hybridizing region as CMPD A and a 10 nucleotide self-complementary region, was mixed with the same 25-mer oligonucleotides at room temperature. The mixture was then gradually heated and increase in hyperchromicity was plotted against increase in temperature. The results are

20

-29-

shown as a solid line in Figure 3. This time, in addition to the melting observed at about 65°C, an earlier increase in hyperchromicity was observed at about 58°C, corresponding to the melting of the intra molecular hydrogen bonds of the hairpin structure. This result indicates that the intramolecular base pairing involving the self-complementary region is less thermodynamically stable than the intermolecular base pairing between the target hybridizing region and a complementary oligonucleotide.

To further test the increased stability of the intermolecular base pairing relative to the intramolecular base pairing, CMPD B was then mixed with the same complementary 25-mer oligonucleotide and heated to 80°C, then allowed to cool to room temperature. This mixture was then gradually heated and increase in hyperchromicity was plotted against increase in temperature. The results are shown as a dashed line in Figure 3. Only a single melting temperature of about 65°C was observed, indicating that the intermolecular base pairing between CMPD B and the complementary 25-mer oligonucleotide is favored in competition with intramolecular base pairing involving the self-complementary region.

These results demonstrate that self-stabilized oligonucleotides will hybridize to complementary nucleic acid sequences notwithstanding the presence of oligonucleotide sequences within the oligonucleotide that are complementary to the target hybridizing region. Since it is well known that certain types of oligonucleotide structures hybridize more stably than certain other types of oligonucleotide structures (e.g., RNA:DNA hybrids > DNA:DNA hybrids and phosphodiester - containing oligos > phosphorothioate methylphosphonate or phosphoramidate - containing oligos), these results also indicate that the preferential target hybridizing effect

-30-

may be enhanced by designing the self-stabilized oligonucleotide such that the hybridization between the target hybridizing region and the target sequence involves more stably pairing oligonucleotide structures than the hybridization involving the self-complementary region.

Those skilled in the art will recognize that self-complementary regions can be prepared according to the above teachings and combined with a wide variety of target hybridizing regions.

EXAMPLE 5**Hyperstabilized Self-Stabilized Oligonucleotides**

To provide oligonucleotides having a more stable interaction between the self-complementary region and the target hybridizing region, oligodeoxynucleoside phosphodiester or oligodeoxynucleoside phosphorothioates were prepared that had 2-O-Me-ribonucleosides in the self-complementary region. As shown in Table V below, such oligonucleotides had a hyperstabilized interaction between the self-complementary region and the target hybridizing region. Nevertheless, these oligonucleotides continued to favor formation of intermolecular hybrids with complementary DNA, relative to molecules containing intramolecular hybrids.

TABLE V

DUPLEX STABILITY OF SELF-STABILIZED OLIGONUCLEOTIDES HAVING 2-O-Me-RIBONUCLEOTIDES IN THE SELF-COMPLEMENTARY REGION

	TM	Complementary with DNA (25 mer)
5' -CTCTGGCACCCATCTCTCTCCTTCTGGAGA-3'	59°C	64.8°C
5' -CTCTGGCACCCATCTCTCTCCTTCTGGAGAG-3'	66°C	64.5°C
5' -CTCTGGCACCCATCTCTCTCCTTCTGGAGAGAG-3'	71°C	65°C

-31-

Another class of hyperstabilized self-stabilized oligonucleotides was prepared by covalently linking an acridine molecule to the terminus of the self-complementary region. These molecules also demonstrated hyperstability of the interaction between the target hybridizing region and the self-complementary region. Nevertheless, these molecules still preferentially formed intermolecular hybrids with complementary DNA, relative to forming intramolecular hybrids.

10

TABLE VI

DUPLEX STABILITY OF SELF-STABILIZED OLIGONUCLEOTIDES HAVING INTERCALATING AGENTS IN THE SELF-COMPLEMENTARY REGION

	TM	Complementary with DNA (25 mer)
5'-CTCTGGCACCAGATCTCTCTCCTTCTX	N/A	67.5°C
5'-CTCTGGCACCAGATCTCTCTCCTTCTGGX-3'	N/A	66.7°C
5'-CTCTGGCACCAGATCTCTCTCCTTCTGGAGX-3'	65°C	66.3°C
5'-CTCTGGGACCAGATCTCTCTCCTTCTGGAGAGX-3'	66.8°C	66.7°C

15

20

These results indicate that it is possible to construct hyperstabilized self-stabilized oligonucleotides having very stable interactions between the self-complementary region and the target hybridizing region, without interfering with the ability of the oligonucleotide to form intermolecular hybrids with a target nucleic acid.

-32-

We claim:

1. A therapeutic self-stabilized oligonucleotide comprising a target hybridizing region and a self-complementary region,
5 wherein the target hybridizing region comprises a oligonucleotide sequence complementary to a nucleic acid sequence that is from a virus, a pathogenic organism or a cellular gene, and wherein the self-complementary region comprises an oligonucleotide sequence
10 complementary to a nucleic acid sequence that is within the therapeutic self-stabilized oligonucleotide.

2. A therapeutic self-stabilized oligonucleotide according to claim 1, wherein the target hybridizing region comprises four or more contiguous
15 deoxyribonucleotide phosphodiester, phosphorothioates, or phosphorodithioates.

3. A therapeutic self-stabilized oligonucleotide according to claim 1, wherein the self-complementary region comprises nucleotides selected from the group
20 consisting of: deoxyribonucleotide or ribonucleotide phosphodiester, phosphotriester phosphorothioates, phosphorodithioates, phosphoramidates, alkylphosphonates, alkylphosphonothioates, ketones, sulfones and sulfates.

4. A therapeutic self-stabilized oligonucleotide
25 according to claim 1, wherein the virus is selected from the group consisting of: human immunodeficiency virus, herpes simplex virus, human papilloma virus, influenza virus, foot and mouth disease virus, yellow fever virus, Varicella-Zoster virus, and cucumber mosaic virus.

-33-

5. A therapeutic self-stabilized oligonucleotide according to claim 1, wherein the pathogenic organism is selected from the group consisting of Plasmodium falciparum, Trypanosoma brucei, Leishmania, and Fasciola hepatica.
5
6. A therapeutic self-stabilized oligonucleotide according to claim 1, wherein the cellular gene is selected from the group consisting of prion protein, Alzheimer's amyloid-like protein, and oncogenes or proto-oncogenes.
10
7. A therapeutic self-stabilized oligonucleotide according to claim 2, wherein the self-complementary region comprises nucleotides selected from the group consisting of: deoxyribonucleotide or ribonucleotide phosphodiester, phosphorothioate, phosphorodithioate, phosphoramidate, alkylphosphonate, alkylphosphonothioate, phosphoromorpholidate, ketone, sulfone and sulfate.
15
8. A therapeutic self-stabilized oligonucleotide according to claim 2, wherein the virus is selected from the group consisting of: human immunodeficiency virus, herpes simplex virus, human papilloma virus, influenza virus, foot and mouth disease virus, yellow fever virus, Varicella-Zoster virus, and cucumber mosaic virus.
20
9. A therapeutic self-stabilized oligonucleotide according to claim 2, wherein the pathogenic organism is selected from the group consisting of Plasmodium falciparum, Trypanosoma brucei, Leishmania, and Fasciola hepatica.
25
10. A therapeutic self-stabilized oligonucleotide according to claim 2, wherein the cellular gene is
30

-34-

selected from the group consisting of prion protein, Alzheimer's amyloid-like protein, and oncogenes or proto-oncogenes.

- 5 11. A therapeutic self-stabilized oligonucleotide according to claim 3, wherein the virus is selected from the group consisting of: human immunodeficiency virus, herpes simplex virus, human papilloma virus, influenza virus, foot and mouth disease virus, yellow fever virus, Varicella-Zoster virus, and cucumber mosaic virus.
- 10 12. A therapeutic self-stabilized oligonucleotide according to claim 3, wherein the pathogenic organism is selected from the group consisting of Plasmodium falciparum, Trypanosoma brucei, Leishmania, and Fasciola hepatica.
- 15 13. A therapeutic self-stabilized oligonucleotide according to claim 3, wherein the cellular gene is selected from the group consisting of prion protein, Alzheimer's amyloid-like protein, and oncogenes or proto-oncogenes.
- 20 14. A self-stabilized ribozyme having a self-complementary region at either or both of its 5' and 3' ends.
- 25 15. A method of inhibiting the gene expression of a virus, a pathogenic organism, or a cellular gene, the method comprising providing the self-stabilized oligonucleotide of claim 1 to virus or pathogen infected cells, or to uninfected cells, respectively.
- 30 16. A method of inhibiting the gene expression of a virus, a pathogenic organism, or a cellular gene, the method comprising providing the self-stabilized

-35-

oligonucleotide of claim 2 to virus or pathogen infected cells, or to uninfected cells, respectively.

17. A method of inhibiting the gene expression of a virus, a pathogenic organism, or a cellular gene, the method comprising providing the self-stabilized oligonucleotide of claim 3 to virus or pathogen infected cells, or to uninfected cells, respectively.

18. A method of treating a diseased human or animal having a disease resulting from a virus or pathogenic organism infection or from the abnormal expression or product of a cellular gene, the method comprising administering to the diseased human or animal oligonucleotide according to claim 1 in a pharmaceutically acceptable carrier.

19. A method of treating a diseased human or animal having a disease resulting from a virus or pathogenic organism infection or from the abnormal expression or product of a cellular gene, the method comprising administering to the diseased human or animal oligonucleotide according to claim 2 in a pharmaceutically acceptable carrier.

20. A method of treating a diseased human or animal having a disease resulting from a virus or pathogenic organism infection or from the abnormal expression or product of a cellular gene, the method comprising administering to the diseased human or animal oligonucleotide according to claim 3 in a pharmaceutically acceptable carrier.

21. A hyperstabilized self-stabilized oligonucleotide having one or more ribonucleotide or 2'-O-Me-ribonucleotide in the self-complementary region or in the

-36-

complementary portion of the target hybridizing region.

22. A hyperstabilized self-stabilized oligonucleotide having one or more intercalating agent molecule in the self-complementary region or in a linker region.

1/7




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 93/06326

A. CLASSIFICATION OF SUBJECT MATTER
IPC 5 C12N15/11 C07H21/00 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 5 C12N C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBO JOURNAL vol. 8, no. 13, 1989, EYNHAM, OXFORD GB pages 4297 - 4305 CASE C.C. ET AL 'The unusual stability of the IS10 anti-sense RNA is critical for its function and is determined by the structure of the stem-domain'	1-21
Y	see page 4303, column 2 ---	22
Y	WO,A,92 03464 (MICROPROBE CORPORATION) 5 March 1992 see claims; example 32 ---	22
	-/--	

Further documents are listed in the continuation of box C. Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search: 16 November 1993

Date of mailing of the international search report: 30. 11. 93

Name and mailing address of the ISA: European Patent Office, P.B. 5818 Patentlaan 2, NL - 2280 HV Rijswijk, Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016

Authorized officer: DAY, G

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 93/06326

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	NUCLEIC ACIDS RESEARCH vol. 21, no. 11 , 11 June 1993 , ARLINGTON, VIRGINIA US pages 2729 - 2735 TANG J.Y. ET AL 'Self-stabilized antisense oligodeoxynucleotide phosphorothioates: properties and anti-HIV activity'	1-21
P,Y	see page 2733, column 2, line 20 - page 2735 -----	22

1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 93/06326

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 18-20 and claims 15-17 (partially) are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 93/06326

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9203464	05-03-92	EP-A- 0547142	23-06-93

This Page Blank (uspto)